arXiv:hep-th/0602048v3 25 Mar 2006

Vacuum Expectation Value of the Spinor Massive field in the Cosmic String Space-Time

Valdir B. Bezerra* and Nail R. Khusnutdinov† ‡
Departamento de Física, Universidade Federal da Paraíba,
Caixa Postal 5008, CEP 58051-970 João Pessoa, Pb, Brazil

We found the contribution to the vacuum expectation value of the energy-momentum tensor of a massive Dirac field due to the conical geometry of the cosmic string space-time. The heat kernel and heat kernel expansion for the squared Dirac operator in this background are also considered and the first three coefficients were found in an explicity form.

I. INTRODUCTION

The study of quantum fields in the space-time of a static cylindrically symmetric cosmic string has been done in different situations. Examples of these studies are the computation of the non-vanishing contribution to the vacuum expectation value of the energy-momentum tensor of quantum fields, as for example, scalar, spinor and vector fields [2, 4, 11, 14, 17, 23, 25, 39, 41, 44, 45, 49]. Quantum fields at non-zero temperature were also considered in the literature [9, 10, 18, 24, 35, 36, 37, 38, 42]. It was noted that the energy-momentum tensor has a non-integrable singularity at the origin and possesses the structure similar to the energy-momentum tensor in the wedge [13]. On the other hand there is a great interest connected with properties of the heat kernel of the Laplacian like operator and its expansion in this background. The point to be considered is that in this space-time there is a singular surface with codimension two which originates some problems [33]. As a consequence, it is impossible to apply standard expressions for the heat kernel coefficients as discussed in [46] and therefore we have to modify them. Nevertheless, there is a way to calculate the heat kernel coefficients by considering generalized cone in dimensions greater then four [7] and to treat the heat kernel coefficients in the standard way using dimensional reduction. General considerations concerning the first heat kernel coefficient and up to systems with spin-2 were considered in Ref. [20].

Cosmic string [31, 47] is an exotic topological defect [48] which may have been formed at phase transitions in the very early history of the Universe. Up to the moment no direct observational evidence of their existence has been found (see, nevertheless [40]), but the richness of the new ideas they brought along to general relativity seems to justify the interest in the study of these structures.

The gravitational field of a straight infinitely long cosmic string is quite remarkable; a particle placed at rest around a straight, infinite, static string will not be attracted to it; there is no local gravity. The space-time around a cosmic string is locally flat but not globally. The external gravitational field due to a cosmic string may be approximately described by a commonly called conical geometry. Due to this conical geometry a cosmic string can induce several effects like, for example, gravitational lens [21, 47], pair production [26], electrostatic self-force [34] on an electric charge at rest, bremsstrahlung process [1, 6] and the so-called gravitational Aharonov-Bohm effect [16].

In the present paper we obtain an explicit form of the expression for vacuum expectation value of the energy-momentum tensor of a massive Dirac field in the cosmic string space-time, showing up that the conical structure of this space-time induces a non-vanishing value for this quantity. The massless case was already considered by Frolov and Serebriany [17]. The energy-momentum tensor is traceless and it has the same non-integrable singularity at the string as in the massless case. We also reobtain the heat kernel and heat kernel expansion in terms of contour integrals. We show that in the coincidence limit, the Green function of squared Dirac operator is defined by the first heat kernel coefficient and the energy-momentum tensor is defined by the second one. The vacuum expectation value of the energy-momentum tensor of a massive scalar field was considered in Refs. [23, 27].

The paper is organized as follows. In Sec. II we find the Green function of the squared Dirac operator and the corresponding heat kernel. In Sec. III we obtain the expansion of the heat kernel and found in manifest form the heat kernel coefficients which are expressed in terms of the delta function and its derivatives. The vacuum expectation value for the Dirac field is obtained in Sec. IV. We conclude with the Sec.V by summarizing the results obtained presenting some remarks. In this paper we adopt the units where $\hbar = c = G = 1$.

---

[*] e-mail: valdir@fisica.ufpb.br
[†] e-mail: nail@kazan-spu.ru
[‡] On leave from Department of Physics, Tatar State University of Humanity and Pedagogic, Mezhlauk 1, Kazan 420021, Russia

II. GREEN FUNCTION AND HEAT KERNEL

To start with let us consider the massive Dirac equation in the Euclidean sector, $\tau = it$, in the space-time generated by a cosmic string. The space-time is described by the following line element [47]

$$ds^2 = d\tau^2 + d\rho^2 + \frac{\rho^2}{\nu^2} d\varphi^2 + dz^2$$

where $\tau, z \in \mathbb{R}, \rho \in \mathbb{R}^+$ and $\varphi \in [0, 2\pi]$. The manifold associated with a cosmic string, $\mathcal{M} = \mathbb{R}^2 \otimes \mathcal{C}^2_\nu$, is a direct product of two the dimensional space $\mathbb{R}^2$ and two dimensional conical space $\mathcal{C}^2_\nu$. The scalar curvature has to be understood as a distribution [43]

$$\mathcal{R} = 4\pi \frac{\nu - 1}{\nu} \frac{\delta^{(2)}(x)}{\sqrt{g}}. \tag{1}$$

The general considerations about the properties of this space-time may be found in [8] while the Green function of the Dirac equation, for a massive field in this background was found in Ref. [37]. We also obtain the Green function in this background, but in different form compared with the one obtained in [37] which is more suitable for our proposal.

According the general considerations concerning the Dirac equation in arbitrary dimensions [15, 22, 32], the Euclidean spinor Green function obeys the following equation

$$\not{D}_m \mathcal{S}(x; x') = -I_4 \frac{\delta^{(4)}(x - x')}{\sqrt{g}},$$

where $\not{D}_m = \gamma^\mu \widetilde{\nabla}_\mu + m$ is the Dirac operator and $\widetilde{\nabla}_\mu = \partial_\mu + \Gamma_\mu$. The Fock-Ivanenko spinor connection

$$\Gamma_\mu = \frac{1}{2} \sigma^{ab} e^\nu_{(a)} \nabla_\mu e_{(b)\nu}$$

is expressed in terms of the vierbein $e_{(b)\nu}$ and the matrix $\sigma^{ab} = \frac{1}{4}[\gamma^{(a)}, \gamma^{(b)}]$. Gamma matrices satisfy the Clifford commutation relation $\{\gamma^{(a)}, \gamma^{(b)}\} = 2\delta^{ab} I_4$ and the generalized Dirac matrices, $\gamma^\mu(x)$, are defined by the relation $\gamma^\mu(x) = e^\mu_{(a)} \gamma^{(a)}$. In what follows we will adopt the standard representation

$$\gamma^{(0)} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \; \gamma^{(k)} = -i \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix},$$

where σ^k are the Pauli matrices. As to the vierbein, we will consider the one used in [12], which are given by the following expressions

$$e^\mu_{(a)} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\varphi & -\frac{\nu}{\rho}\sin\varphi & 0 \\ 0 & \sin\varphi & +\frac{\nu}{\rho}\cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

where rows correspond to the number (a) of the vector $e^\mu_{(a)}$. Thus, we obtain the results

$$\gamma^\tau = \gamma^{(0)}, \; \gamma^\rho = \cos\varphi \gamma^{(1)} + \sin\varphi \gamma^{(2)}, \; \gamma^\varphi = -\frac{\nu}{\rho} \sin\varphi \gamma^{(1)} + \frac{\nu}{\rho} \cos\varphi \gamma^{(2)}, \; \gamma^z = \gamma^{(3)},$$

and

$$\Gamma_\mu = -\frac{\nu - 1}{2\nu} \gamma^{(2)} \gamma^{(1)} \delta_{\mu,\varphi} = i \frac{\nu - 1}{2\nu} \Sigma_3 \delta_{\mu,\varphi},$$

where $\Sigma_3 = -i\gamma^1\gamma^2$. Therefore, the spinor Green function $\mathcal{S}$ obeys the following equation

$$(\gamma^\mu \partial_\mu - \frac{\nu - 1}{2\rho} \gamma^\rho + m) \mathcal{S}(x; x') = -I_4 \frac{\delta^4(x - x')}{\sqrt{g}}.$$

Let us define the Green function $\mathcal{G}$ of the squared Dirac operator by the relation

$$\mathcal{S}(x; x') = \not{D}_{-m} \mathcal{G}(x; x'). \tag{2}$$

It obeys the following equation

$$\not{D}^2\mathcal{G}(x;x') = -I_4\frac{\delta^4(x-x')}{\sqrt{g}},$$

where

$$\not{D}^2 = \not{D}_m\not{D}_{-m} = \left(g^{\mu\nu}\partial^2_{\mu\nu} + \frac{1}{\rho}\partial_\rho - \frac{(\nu-1)^2}{4\rho^2} + i\frac{\nu(\nu-1)}{\rho^2}\Sigma_3\partial_\varphi - m^2 - \frac{1}{4}\mathcal{R}\right). \tag{3}$$

This equation has no the form of the equation for the scalar Green function. Its general form was presented in [46]. In this paper we will omit the term with the singular scalar curvature. The contribution arising from it has been discussed in the literature[3, 4, 5, 29].

Due to the fact that Σ_3 is diagonal, the Green function $\mathcal{G}$ is diagonal, too. In order to obtain the Green function in explicitly form let us find the full set of bispinors which satisfies the equation

$$\not{D}^2\phi(x) = -\lambda^2\phi(x). \tag{4}$$

We shall numerate the eigenfunctions by eigenvalues of operators $\widehat{p}_\tau$, $\widehat{p}_z$ and the projection of the full momentum $\widehat{J}_3$, as

$$\begin{aligned}
\widehat{p}_\tau\phi &= -i\partial_\tau\phi = E\phi,\\
\widehat{p}_z\phi &= -i\partial_z\phi = p_3\phi,\\
\widehat{J}_3\phi &= (\widehat{p}_\varphi + \frac{1}{2\nu}\Sigma_3)\phi = (-i\partial_\varphi + \frac{1}{2}\Sigma_3)\phi = j\phi,
\end{aligned}$$

and define the number $p_\perp$ by relation $\lambda^2 = p_\perp^2 + E^2 + p_3^2 + m^2$. All these operators commute with each other and with Dirac and squared Dirac operators. The above set of operators define the eigenfunction up to four arbitrary constants. Let us numerate these independent solutions by eigenvalues of matrices Σ_3 with eigenvalue $a = \pm1$ and $\gamma_5 = \gamma^{(0)}\gamma^{(1)}\gamma^{(2)}\gamma^{(3)}$ with eigenvalue $b = \pm1$. Both these operators commute with the above set of operators. We shall numerate eigenfunctions by numbers $q = E, p_3, p_\perp, l$ where $E, p_3 \in \mathbb{R}, p_\perp \in \mathbb{R}^+, l = 0, \pm1, \ldots (j = l + \frac{1}{2})$ and by $(a,b) = (\pm1,\pm1)$. Therefore we have the following four independent solutions

$$\begin{aligned}
\phi_q^{(+,\pm)} &= \frac{\sqrt{\nu}}{4\pi^{3/2}}e^{iE\tau+ip_3z+il\varphi}J_{\beta_-}(p_\perp\rho)w_{(+,\pm)},\\
\phi_q^{(-,\pm)} &= \frac{\sqrt{\nu}}{4\pi^{3/2}}e^{iE\tau+ip_3z+i(l+1)\varphi}J_{\beta_+}(p_\perp\rho)w_{(-,\pm)},
\end{aligned}$$

where

$$w_{(+,\pm)} = \begin{pmatrix}1\\0\\\pm1\\0\end{pmatrix},\ w_{(-,\pm)} = \begin{pmatrix}0\\1\\0\\\pm1\end{pmatrix}$$

are the eigenfunctions of Σ_3 and γ_5. Here $J_\mu(z)$ is the Bessel function with indices $\beta_\mp = |\nu j \mp \frac{1}{2}|$. This full set obeys the following relations of completeness and orthogonality

$$\sum_l \iiint dEdp_3p_\perp dp_\perp \phi_q^{(a,b)\dagger}(x')\phi_q^{(a,b)}(x) = \frac{\delta^{(4)}(x-x')}{\sqrt{g}},$$

$$\int \phi_{q'}^{(a',b')\dagger}(x')\phi_q^{(a,b)}(x)\sqrt{g}d^4x = \delta_{a,a'}\delta_{b,b'}\delta_{l,l'}\delta(E-E')\delta(p_3-p_3')\frac{\delta(p_\perp-p_\perp')}{p_\perp}.$$

Now we are ready to obtain the Green function in manifest form. To do this, let us represent the Green function in the following form

$$\begin{aligned}
\mathcal{G}(x;x') &= \sum_{a,b}\sum_l \iiint dEdp_3p_\perp dp_\perp\frac{\phi_q^{(a,b)}(x)\otimes\phi_q^{(a,b)\dagger}(x')}{p_\perp^2+E^2+p_3^2+m^2} \qquad (5)\\
&= \frac{\nu}{(2\pi)^3}\sum_l \iiint dEdp_3p_\perp dp_\perp\frac{e^{iE\triangle\tau+ip_3\triangle z}}{p_\perp^2+E^2+p_3^2+m^2}
\end{aligned}$$

$$\times\ \mathrm{diag}(J_{\beta_-}(r)J_{\beta_-}(r')e^{il\triangle\varphi},\ J_{\beta_+}(r)J_{\beta_+}(r')e^{i(l+1)\triangle\varphi},\ J_{\beta_-}(r)J_{\beta_-}(r')e^{il\triangle\varphi},\ J_{\beta_+}(r)J_{\beta_+}(r')e^{i(l+1)\triangle\varphi}),$$

where $r = p_\perp\rho,\ r' = p_\perp\rho'$. By changing the variable $l \to l-1$ in the second and forth terms, we can write the Green function in more simple form, as

$$\begin{aligned}\mathcal{G}(x;x') &= \frac{\nu}{(2\pi)^3}\sum_l \iiint dEdp_3p_\perp dp_\perp \frac{e^{iE\triangle\tau+ip_3\triangle z+il\triangle\varphi}}{p_\perp^2+E^2+p_3^2+m^2}\\ &\times\ \mathrm{diag}(J_{\xi_+}(r)J_{\xi_+}(r'),\ J_{\xi_-}(r)J_{\xi_-}(r'),\ J_{\xi_+}(r)J_{\xi_+}(r'),\ J_{\xi_-}(r)J_{\xi_-}(r')),\end{aligned}$$

where $\xi_\pm = |\nu l \pm \frac{\nu-1}{2}|$.

It is easy to find from this expression the heat kernel of the squared Dirac operator in Eq. (3). It is given by

$$\begin{aligned}\mathcal{K}(x;x'|s) &= \frac{\nu}{(2\pi)^3}\sum_l \iiint dEdp_3p_\perp dp_\perp e^{iE\triangle\tau+ip_3\triangle z+il\triangle\varphi-s(p_\perp^2+E^2+p_3^2+m^2)}\\ &\times\ \mathrm{diag}(J_{\xi_+}(r)J_{\xi_+}(r'),\ J_{\xi_-}(r)J_{\xi_-}(r'),\ J_{\xi_+}(r)J_{\xi_+}(r'),\ J_{\xi_-}(r)J_{\xi_-}(r'))\\ &= \mathcal{K}_m^{(2)}(\triangle\tau,\triangle z|s)\mathcal{K}_\nu^{(2)}(\triangle\varphi,\rho,\rho'|s),\end{aligned}\tag{6}$$

where $\mathcal{K}_m^{(2)}(\triangle\tau,\triangle z|s)$ is the heat kernel of the operator $\widehat{L}_m = \partial^2_{\tau^2}+\partial^2_{z^2}-m^2$ and $\mathcal{K}_\nu^{(2)}(\triangle\varphi,\rho,\rho'|s)$ is the heat kernel of the operator

$$\widehat{L}_\nu = \partial^2_{\rho^2}+\frac{1}{\rho}\partial_\rho+\frac{\nu^2}{\rho^2}\partial^2_{\varphi^2}-\frac{(\nu-1)^2}{4\rho^2}+i\frac{\nu(\nu-1)}{\rho^2}\Sigma_3\partial_\varphi.$$

Integrating over $E, p_3, p_\perp$, we get

$$\mathcal{K}_m^{(2)}(\triangle\tau,\triangle z|s) = \frac{1}{4\pi s}e^{-m^2s-\frac{\triangle\tau^2+\triangle z^2}{4s}},\tag{7a}$$

$$\begin{aligned}\mathcal{K}_\nu^{(2)}(\triangle\varphi,\rho,\rho'|s) &= \frac{\nu}{4\pi s}\sum_l e^{il\triangle\varphi-\frac{\rho^2+\rho'^2}{4s}}\mathrm{diag}(I_{\beta_-}(\frac{\rho\rho'}{2s}),\ I_{\beta_+}(\frac{\rho\rho'}{2s})e^{i\triangle\varphi},\ I_{\beta_-}(\frac{\rho\rho'}{2s}),\ I_{\beta_+}(\frac{\rho\rho'}{2s})e^{i\triangle\varphi})\\ &= \frac{\nu}{4\pi s}\sum_l e^{il\triangle\varphi-\frac{\rho^2+\rho'^2}{4s}}\mathrm{diag}(I_{\xi_+}(\frac{\rho\rho'}{2s}),\ I_{\xi_-}(\frac{\rho\rho'}{2s}),\ I_{\xi_+}(\frac{\rho\rho'}{2s}),\ I_{\xi_-}(\frac{\rho\rho'}{2s})).\end{aligned}\tag{7b}$$

It is worth noting that this expression leads to a heat kernel which is a diagonal matrix. The components of this matrix are defined in terms of the first component $\mathcal{K}_{1,1}$ which we denote by $\mathbf{K}$ by relations: $\mathcal{K}_{3,3} = \mathbf{K}$ and $\mathcal{K}_{2,2} = \mathcal{K}_{4,4} = \mathbf{K}^*$. Indeed, because $\xi_\pm \to \xi_\mp$ with changing $l \to l$ then

$$\sum_l e^{il\triangle\varphi}I_{\xi_-} = \sum_l e^{-il\triangle\varphi}I_{\xi_+} = \left(\sum_l e^{il\triangle\varphi}I_{\xi_+}\right)^*,$$

and the heat kernel has the following structure

$$\mathcal{K}_\nu^{(2)} = \mathrm{diag}\left(\mathbf{K},\mathbf{K}^*,\mathbf{K},\mathbf{K}^*\right).\tag{7c}$$

Using Eqs. (2) and (6) we obtain the spinor Green function in manifest form, which can be written as

$$\mathcal{S}(x;x') = \frac{\nu}{(4\pi)^2}\sum_l e^{il\triangle\varphi}\int_0^\infty \frac{ds}{s^2}e^{-sm^2-\frac{\triangle\tau^2+\triangle z^2+\rho^2+\rho'^2}{4s}}(-mS_1-\frac{\triangle\tau}{2s}S_2+i\frac{\triangle\ z}{2s}S_3-\frac{i}{2s}S_4)\tag{8}$$

where

$$\begin{aligned}S_1 &= \mathrm{diag}\left(I_{\beta_-}, I_{\beta_+}e^{i\triangle\varphi}, I_{\beta_-}, I_{\beta_+}e^{i\triangle\varphi}\right),\ S_2 = \mathrm{diag}\left(I_{\beta_-}, I_{\beta_+}e^{i\triangle\varphi}, -I_{\beta_-}, -I_{\beta_+}e^{i\triangle\varphi}\right),\\ S_3 &= \begin{pmatrix} 0 & 0 & I_{\beta_-} & 0\\ 0 & 0 & 0 & -I_{\beta_+}e^{i\triangle\varphi}\\ -I_{\beta_-} & 0 & 0 & 0\\ 0 & I_{\beta_+}e^{i\triangle\varphi} & 0 & 0\end{pmatrix},\\ S_4 &= \begin{pmatrix} 0 & 0 & 0 & e^{-i\varphi'}(\rho' I_{\beta_-}-\rho I_{\beta_+})\\ 0 & 0 & e^{i\varphi}(\rho' I_{\beta_+}-\rho I_{\beta_-}) & 0\\ 0 & -e^{-i\varphi'}(\rho' I_{\beta_-}-\rho I_{\beta_+}) & 0 & 0\\ -e^{i\varphi}(\rho' I_{\beta_+}-\rho I_{\beta_-}) & 0 & 0 & 0\end{pmatrix}\end{aligned}\tag{9}$$

and the argument of the Bessel functions of second kind $I_{\beta_\pm}$ is $z = \frac{\rho\rho'}{2s}$. To obtain previous expression we used the relations :

$$I'_{\beta_-} - \frac{\epsilon_l \beta_-}{z} I_{\beta_-} = I_{\beta_+},\ I'_{\beta_+} + \frac{\epsilon_l \beta_+}{z} I_{\beta_+} = I_{\beta_-},$$

where $\epsilon_l = \mathrm{sgn}(l)$.

Now, let us consider some limiting cases. In the massless case, $m = 0$, the Green function can be found in a closed form and is given explicitly by

$$\mathcal{G}(x;x') = \frac{\nu}{8\pi^2} \frac{1}{\rho\rho'} \frac{1}{\sinh\eta} \frac{\sinh(\frac{\nu+1}{2}\eta) I_4 + \sinh(\frac{\nu-1}{2}\eta) e^{-i\triangle\varphi\Sigma_3}}{\cosh\nu\eta - \cos\triangle\varphi}.$$

We note that this Green function does not coincide with that obtained in [17, 37] and it does not obey the relation

$$\mathcal{G}(\varphi + 2\pi) = -\mathcal{G}(\varphi).$$

Instead of this, it obeys the relation above with minus sign changed to plus sign. The reason for this fact is connected with the use of another vierbein as compared with the one used in [17].

In order to reobtain the result showed in [17], we have to apply a rotation of the vierbein we have chosen by an angle φ, in the plane (ρ, φ). It is well-known that the bispinor will be multiplied by a factor $e^{\frac{i}{2}\varphi\Sigma_3}$. The Green function, as a bispinor at point x and point x', is multiplied by factor $e^{\frac{i}{2}\triangle\varphi\Sigma_3}$. Thus, multiplying by this factor we arrive at the formula

$$\mathcal{G}(x;x') = \frac{\nu}{8\pi^2} \frac{1}{\rho\rho'} \frac{1}{\sinh\eta} \frac{e^{\frac{i}{2}\triangle\varphi\Sigma_3} \sinh(\frac{\nu+1}{2}\eta) + e^{-\frac{i}{2}\triangle\varphi\Sigma_3} \sinh(\frac{\nu-1}{2}\eta)}{\cosh\nu\eta - \cos\triangle\varphi}$$

which coincides with the results obtained in [17, 37], if we take into account that $i\Sigma_3 = \gamma^{(1)}\gamma^{(2)}$.

In the case of zero angle deficit, $\nu = 1$, we have $\xi_\pm = |l|$ and thus

$$\mathcal{K}(x;x'|s) = \frac{1}{(4\pi s)^2} e^{-m^2 s - \frac{R^2}{4s}} I_4,$$

where $R = (\triangle\tau^2 + \triangle z^2 + \rho^2 + \rho'^2 - 2\rho\rho'\cos\triangle\varphi)^{1/2}$ is the distance between points x and x' in the Euclidean 4-dimensional space in cylindrical coordinates. The Green function in this case turns into

$$\mathcal{G}(x;x') = \int_0^\infty \mathcal{K}(x;x'|s) ds = \frac{1}{4\pi^2} \frac{m}{R} K_1(mR) I_4,$$

as it should be. Here K_μ is the Bessel function of second kind.

III. HEAT KERNEL EXPANSION

Let us investigate the asymptotic expansion of the heat kernel of the squared Dirac operator which is given by Eqs. (7b) and (7c). It is impossible to take an expansion of the Bessel function because the asymptotic expansion of the Bessel function depends on the ratio argument and indices. For this reason we use the integral representation for the Bessel function, as follows

$$I_\mu(z) = \frac{1}{2\pi} \int_\Gamma e^{i(w - \frac{\pi}{2})\mu + z\sin w} dw.$$

The contour Γ lies in the half-strip $\Re z \in [-\frac{\pi}{2}, \frac{3\pi}{2}], \Im z > 0$. It goes from $-\frac{\pi}{2} + i\infty$ to $\frac{3\pi}{2} + i\infty$.

Let us consider, in general, the series

$$f(\alpha, r) = \nu \sum_l e^{il\triangle\varphi} I_{|\nu(l+\alpha)|}(r),$$

where $|\alpha| \leq 1$. For our cases we have $\alpha = \pm\frac{\nu-1}{2\nu}$ and $|\alpha| < \frac{1}{2}$. Then we use the representation above for the Bessel function and change the integration and summation and shift the variable w to $z = w - \frac{\pi}{2}$. In this situation, we obtain the following expression for the function $f(\alpha, r)$

$$f(\alpha, r) = \frac{\nu}{2\pi} \int_{\Gamma'} e^{r\cos z} \left[\frac{e^{-iz|\nu\alpha|}}{e^{-i(z\nu - \epsilon_\alpha\triangle\varphi)} - 1} - \frac{e^{iz|\nu\alpha|}}{e^{i(z\nu + \epsilon_\alpha\triangle\varphi)} - 1} \right] dz,$$

where $\epsilon_\alpha = sgn(\alpha)$. The contour Γ' is the contour Γ shifted by $\pi/2$ to the left. It lies in the positive part of the strip $z \in [-\pi, \pi]$. Then we divide the integral in two parts according with integrand and in the second part, we do the change of variable $z \to -z$. The integrand takes the form as the first integrand but the contour is the central symmetry of Γ'. Therefore we may recombine both integrals into a single integral and arrive at the following formula for function $f(\alpha, r)$

$$f(\alpha, r) = \frac{\nu}{2\pi} \int_\gamma \frac{e^{-iz|\nu\alpha| + r\cos z}}{e^{-i(z\nu - \epsilon_\alpha \triangle\varphi)} - 1} dz.$$

The contour γ has two branches. First one is contour Γ' and the second is the central symmetry of Γ'. Taking into account this formula we arrive at the following expressions for heat kernel component

$$\mathbf{K} = \frac{e^{-\frac{\rho^2 + \rho'^2}{4s}}}{4\pi s} f(\frac{\nu - 1}{2\nu}, \frac{\rho\rho'}{2s}) = \frac{\nu e^{-\frac{\rho^2 + \rho'^2}{4s}}}{8\pi^2 s} \int_\gamma \frac{e^{-iz\frac{\nu-1}{2} + \frac{\rho\rho'}{2s}\cos z}}{e^{-i(z\nu - \triangle\varphi)} - 1} dz.$$

The complex conjugate $\mathbf{K}^*$ differ from $\mathbf{K}$ by the sign of $\triangle\varphi$ only.

Now, let us return to the function $f(\alpha, r)$ and modify the contour γ. It is easy to see that the zeros of the denominator are in the points $z_n = \epsilon_\alpha \frac{\triangle\varphi}{\nu} + \frac{2\pi n}{\nu}$. Because of $|\triangle\varphi| < \pi$ and $\nu > 1$, then we have that$|\triangle\varphi/\nu| < \pi$. Therefore, the first zero of the denominator, $z_0 = \epsilon_\alpha \frac{\triangle\varphi}{\nu}$, belongs to the interval of integration $[-\pi, \pi]$. Let us extract this first pole in manifest form. With this we add the integrals over two lines around point $z = \epsilon_\alpha \triangle\varphi/\nu$ in each of which we integrate in opposite directions. The contribution from these additional integrals are zero. Then we divide the contour γ with additional lines in two parts and represent the integral as an integral over contour χ and over a circle around point $\epsilon_\alpha \frac{\triangle\varphi}{\nu}$. The contour χ has two branches. The first one goes from $-\pi + i\infty$ to $-\pi - i\infty$, very close to the point $\epsilon_\alpha \frac{\triangle\varphi}{\nu}$ from the left side. The second part of the contour goes from $\pi + i\infty$ to $\pi - i\infty$, very close to the point $\epsilon_\alpha \frac{\triangle\varphi}{\nu}$ from the right side. The second integral is minus the residue at point $\epsilon_\alpha \frac{\triangle\varphi}{\nu}$. Therefore we may represent our function $f(\alpha, r)$ in the form below

$$f(\alpha, r) = e^{r\cos\frac{\triangle\varphi}{\nu} - i\alpha\triangle\varphi} + \frac{\nu}{2\pi} \int_\chi \frac{e^{-iz|\nu\alpha| + r\cos z}}{e^{-i(z\nu - \epsilon_\alpha \triangle\varphi)} - 1} dz.$$

Taking into account this representation we arrive at the following expression for component $\mathbf{K}$

$$\mathbf{K} = \frac{e^{-\frac{d^2}{4s} - i\frac{\nu-1}{2\nu}\triangle\varphi}}{4\pi s} \left[1 + \frac{\nu}{2\pi} \int_\chi \frac{e^{-i(z\nu - \triangle\varphi)\frac{\nu-1}{2\nu} + \frac{\rho\rho'}{2s}(\cos z - \cos\frac{\triangle\varphi}{\nu})}}{e^{-i(z\nu - \triangle\varphi)} - 1} dz \right],$$

where $d^2 = \rho^2 + \rho'^2 - 2\rho\rho' \cos\frac{\triangle\varphi}{\nu}$. The expression for the complex conjugate $\mathbf{K}^*$ component differs from the above by the sign of $\triangle\varphi$.

Therefore, in the coincidence limit $\rho' = \rho$ and $\varphi' = \varphi$ we obtain the following expression for the heat kernel

$$\mathcal{K}^{(2)}_\nu(x; x|s) = \frac{1}{4\pi s} \left[1 + \frac{\nu}{2\pi} \int_\chi \frac{e^{-iz\frac{\nu-1}{2} - \frac{\rho^2}{s}\sin^2\frac{z}{2}}}{e^{-iz\nu} - 1} dz \right] I_4, \tag{10}$$

where the contour χ intersects the real axis close to the origin. In order to find the expansion of the heat kernel over s we use the formula presented in [9], which is given by

$$\frac{\nu\sigma}{\pi} e^{-\sigma\rho^2} = \sum_{n=0}^{\infty} \frac{\triangle^n_{(2)} \delta^{(2)}(\vec{r})}{n!(4\sigma)^n},$$

where $\delta^{(2)}(\vec{r}) = \frac{\nu}{\rho}\delta(\rho)\delta(\varphi)$ and $\triangle_{(2)} = \partial^2_{\rho^2} + \frac{1}{\rho}\partial_\rho + \frac{\nu^2}{\rho^2}\partial^2_{\varphi^2}$. The integral may be calculated by the method of residue at the zero point. Thus, we obtain

$$\mathcal{K}^{(2)}_\nu(x; x|s) = \frac{1}{4\pi s} \left[I_4 + \sum_{n=1}^{\infty} a_n(x; x) s^n \right],$$

where

$$a_n(x; x) = I_4 \frac{R_{n-1}}{(n-1)! 4^{n-1}} \triangle^{n-1}_{(2)} \delta^{(2)}(\vec{r})$$

and

$$R_n = \mathrm{Res}_0 \frac{i\pi e^{-iz\frac{\nu-1}{2}}}{(e^{-iz\nu}-1)\sin^{2n+2}\frac{z}{2}}.$$

Therefore, the first three coefficients are given by

$$R_0 = -\frac{\pi(\nu^2-1)}{6\nu},\ R_1 = \frac{7\nu^2+17}{60}R_0,\ R_2 = \frac{31\nu^4+178\nu^2+367}{2520}R_0.$$

The expansion of the four dimensional heat kernel is expressed in terms of the same coefficients, as

$$\mathcal{K}(x;x'|s) = \frac{1}{(4\pi s)^2}\left[I_4 + \sum_{n=1}^{\infty} a_n(x;x)s^n\right].$$

From the above expression we observe that the first coefficient R_0 for the spinor field is minus 1/2 of the same coefficient corresponding to the scalar field. Therefore, we have

$$\mathrm{tr} a_1^{(1/2)} = -\frac{N}{2}a_1^{(0)} = -2a_1^{(0)},$$

and thus in accordance with [19, 20, 28]. The other coefficients have different structure.

IV. VACUUM EXPECTATION VALUES

First of all let us find in closed form the coincidence limit of the renormalized Green function of the squared Dirac operator. To do the process of renormalization we subtract the same function with $\nu = 1$. Because the second integral in Eq. (10) is zero for $\nu = 1$, we have the following expression in the coincidence limit

$$\mathcal{G}^{ren}(x;x) = \frac{\nu}{32\pi^3}\int_0^{\infty}\frac{ds}{s^2}\int_{\chi}\frac{e^{-iz\frac{\nu-1}{2}-\frac{\rho^2}{s}\sin^2\frac{z}{2}-m^2 s}}{e^{-iz\nu}-1}dz I_4. \tag{11}$$

Firstly, let us consider the massless case. We may integrate easily with respect to s and obtain

$$\mathcal{G}^{ren}(x;x) = \frac{\nu}{32\pi^3\rho^2}\int_{\chi}\frac{e^{-iz\frac{\nu-1}{2}}}{(e^{-iz\nu}-1)\sin^2\frac{z}{2}}dz I_4.$$

Then we close the contour to infinities and calculate this expression with the residue at the point zero. Thus, we get the result

$$\mathcal{G}^{ren}(x;x) = \frac{\nu R_0}{16\pi^3\rho^2}I_4 = -\frac{\nu^2-1}{96\pi^2\rho^2}I_4. \tag{12}$$

In the massive case, we deform the contour in two straight lines over $z = \pm\pi$. We have to take into account the residues at points $z_n = 2\pi n/\nu$, where $|n| < \nu/2$. For $\nu < 2$ there is no zeros in the denominator and no residues. By changing the variable $z = \pm\pi + iy$ we obtain, in general, the result

$$\int_{\chi}\frac{e^{-iz\frac{\nu\pm 1}{2}-\frac{\rho^2}{s}\sin^2\frac{z}{2}}}{e^{-iz\nu}-1}dz = 4\cos\frac{\pi\nu}{2}\int_0^{\infty}e^{-\frac{\rho^2}{s}\cosh^2\frac{y}{2}}\frac{\sinh\frac{y}{2}\sinh\frac{y\nu}{2}}{\cosh\nu y-\cos\pi\nu}dy.$$

Taking into account this formula we arrive at the following expression

$$\mathcal{G}^{ren}(x;x) = \frac{m}{16\pi^2\rho}\sum_{n=1}^{[\nu/2]}(-1)^n\tan\frac{\pi n}{\nu}K_1(2m\rho\sin\frac{\pi n}{\nu}) + \frac{m\nu\cos\frac{\pi\nu}{2}}{4\pi^3\rho}\int_0^{\infty}\frac{K_1(2m\rho\cosh\frac{y}{2})}{\cosh\frac{y}{2}}\frac{\sinh\frac{y}{2}\sinh\frac{y\nu}{2}}{\cosh\nu y-\cos\pi\nu}dy I_4.$$

In the case $\nu < 2$, only the last term survives. The appearance of additional terms, when $\nu > 2$, is related with the fact that additional closed geodesics in $\mathcal{C}_{\nu}^2$, appears in this case [30]. For $m\rho \gg 1$ the above expression exponentially falls down, according to

$$\mathcal{G}^{ren}(x;x) \approx \frac{\nu R_0}{16\pi^{5/2}}\frac{m^2 e^{-2m\rho}}{(m\rho)^{3/2}}.$$



FIG. 1: The plot of the ratio $F = \mathcal{G}^{ren}(x;x)_{m\neq0}/\mathcal{G}^{ren}(x;x)_{m=0}$ for $\nu = 0.01$ (thin curve) and $\nu = 2$ (thick curve).

In the opposite case, if $m\rho \ll 1$, the expression for $\mathcal{G}^{ren}(x;x)$ is given by Eq. (12). The plot of the ratio $F = \mathcal{G}^{ren}(x;x)_{m\neq0}/\mathcal{G}^{ren}(x;x)_{m=0}$ is reproduced in Fig. 1 as a function of $m\rho$.

To calculate the energy-momentum tensor we use the following formula [22]

$$\langle T_{\mu\nu}\rangle^{ren} = -\frac{1}{4}\lim_{x'\to x}\mathrm{Im}\{\mathrm{Tr}[\gamma^L_{(\mu}(\nabla_{\nu)}[\mathcal{S}+\mathcal{S}^c]^{ren} - g_{\nu)}^{\cdot\lambda'}\nabla_{\lambda'}[\mathcal{S}+\mathcal{S}^c]^{ren})I(x';x)]\}, \tag{13}$$

where $\mathcal{S}^c$ is the charge conjugate spinor Green function $\mathcal{S}^c = -C\gamma_L^{(0)T}\mathcal{S}^*\gamma_L^{(0)}C^\dagger = -\gamma_L^{(2)}\mathcal{S}^*\gamma_L^{(2)}$. In above formulas γ_L^μ are the gamma matrices in Lorentz regime: $\gamma^{(0)} = \gamma_L^{(0)}$, $\gamma^{(k)} = -i\gamma_L^{(k)}$, and $\tau = it$. In manifest form

$$\mathcal{S}^c(x;x') = \frac{\nu}{(4\pi)^2}\sum_l e^{-il\triangle\varphi}\int_0^\infty \frac{ds}{s^2}e^{-sm^2-\frac{\triangle\tau^2+\triangle z^2+\rho^2+\rho'^2}{4s}}(-mS_1^c - \frac{\triangle\tau^*}{2s}S_2^c + i\frac{\triangle\ z}{2s}S_3^c - \frac{i}{2s}S_4^c) \tag{14}$$

where

$$\begin{aligned}
S_1^c &= \mathrm{diag}\left(I_{\beta_+}e^{-i\triangle\varphi}, I_{\beta_-}, I_{\beta_+}e^{-i\triangle\varphi}, I_{\beta_-}\right),\ S_2^c = \mathrm{diag}\left(-I_{\beta_+}e^{-i\triangle\varphi}, -I_{\beta_-}, I_{\beta_+}e^{-i\triangle\varphi}, I_{\beta_-}\right),\\
S_3^c &= \begin{pmatrix} 0 & 0 & I_{\beta_+}e^{-i\triangle\varphi} & 0\\ 0 & 0 & 0 & -I_{\beta_-}\\ -I_{\beta_+}e^{-i\triangle\varphi} & 0 & 0 & 0\\ 0 & I_{\beta_-} & 0 & 0\end{pmatrix},\\
S_4^c &= \begin{pmatrix} 0 & 0 & 0 & e^{-i\varphi}(\rho' I_{\beta_+} - \rho I_{\beta_-})\\ 0 & 0 & e^{i\varphi'}(\rho' I_{\beta_-} - \rho I_{\beta_+}) & 0\\ 0 & -e^{-i\varphi}(\rho' I_{\beta_+} - \rho I_{\beta_-}) & 0 & 0\\ -e^{i\varphi'}(\rho' I_{\beta_-} - \rho I_{\beta_+}) & 0 & 0 & 0\end{pmatrix}
\end{aligned} \tag{15}$$

and the argument of the Bessel functions is $z = \frac{\rho\rho'}{2s}$. The manifest form of the spinor Green function $\mathcal{S}$ is given by Eq. (8). In order to renormalize the spinor Green functions $\mathcal{S}$ and $\mathcal{S}^c$ we have to subtract from them the same expression with $\nu = 1$. It means that the renormalized Green functions have the same form in which we have to make the changes

$$\nu I_{\beta_-} \to \nu I_{\beta_-} - I_{|l|},\ \nu I_{\beta_+} \to \nu I_{\beta_+} - I_{|l+1|}.$$

In what follows we assume these changes in the Green functions. Because the renormalized Green functions are finite in the coincidence limit we may use $\delta_\nu^{\lambda'}$ instead of $g_\nu^{\cdot\lambda'}$. To obtain the energy-momentum tensor we use the following relations

$$\sum_l I_{\beta_-} = \sum_l I_{\beta_+}, \tag{16a}$$

$$\sum_l l I_{\beta_-} = -\sum_l (l+1) I_{\beta_+}, \tag{16b}$$

which can be easy proved by changing $l \to -l-1$ in the sum in the left hand side. Let us briefly show the main steps of calculations (for simplicity we consider the case $\nu < 2$).

From Eq. (13) we conclude that the energy-momentum tensor has diagonal form with components (no summation):

$$T^\mu_\mu = -\frac{1}{4}\lim_{x'^\mu \to x^\mu} \mathrm{Im}\ \mathrm{Tr}\gamma^\mu_L(\partial_\mu - \partial_{\mu'})(\mathcal{S} + \mathcal{S}^c)^{ren}.$$

In this formula the coincidence limit for all variables except x^μ has already done. The contributions from $\mathcal{S}^c$ coincides with contribution from $\mathcal{S}$. Let us consider consequently different components of the energy-momentum tensor.

T^t_t: There is non-zero contribution only due to S_2, because S_1 and S_4 do not depend on the time:

$$T^t_t = \frac{1}{8\pi^2}\int_0^\infty \frac{ds}{s^3} e^{-sm^2}\left\{e^{-\frac{\rho^2}{2s}}\nu\sum_l I_{\beta_-}(\frac{\rho^2}{2s})\right\}^{ren}.$$

T^z_z: The calculations of the T^z_z component is similar. The non-zero contribution is only due to S_3 and we obtain $T^z_z = T^t_t$, as should be the case due to boost-invariance along z-axis.

T^ρ_ρ: In the coincidence limit S_1 and S_4 survive. The term S_1 gives no contribution because

$$\lim_{\rho' \to \rho}(\partial_\rho - \partial_{\rho'})e^{-\frac{\rho^2+\rho'^2}{4s}} I_{\beta_\pm}(\frac{\rho\rho'}{2s}) = 0.$$

The non-zero contribution appears only due to S_4. By using the relations

$$\lim_{\rho' \to \rho}(\partial_\rho - \partial_{\rho'})\left\{\rho' I_{\beta_\mp}(\frac{\rho\rho'}{2s}) - \rho I_{\beta_\pm}(\frac{\rho\rho'}{2s})\right\} \;=\; -I_{\beta_-}(\frac{\rho^2}{2s}) - I_{\beta_+}(\frac{\rho^2}{2s})$$

and (16a) one has

$$\lim_{\rho' \to \rho}\gamma^\rho_L(\partial_\rho - \partial_{\rho'})S_4 = -2I_{\beta_-}(\frac{\rho^2}{2s})\gamma^{\rho\,2}_L = 2I_{\beta_-}(\frac{\rho^2}{2s})I_4.$$

Therefore, we obtain $T^\rho_\rho = T^t_t$.

T^φ_φ: In the coincidence limit S_1 and S_4 survive. The component S_1 has no contribution to T^φ_φ. Indeed, by using Eq. (16b) one has

$$\mathrm{Tr}\lim_{\varphi' \to \varphi}\gamma^\varphi_L(\partial_\varphi - \partial_{\varphi'})e^{il\triangle\varphi}S_1 = 2ilI_{\beta_-}(\frac{\rho^2}{2s})\mathrm{Tr}\gamma^\varphi_L \mathrm{diag}(1,-1,1,-1) = 0.$$

The non-zero contribution exists only due to S_4. One has

$$\begin{aligned}\lim_{\varphi' \to \varphi}(\partial_\varphi - \partial_{\varphi'})e^{il\triangle\varphi}S_4 \;&=\; 2il\lim_{\varphi' \to \varphi}S_4 + \lim_{\varphi' \to \varphi}(\partial_\varphi - \partial_{\varphi'})S_4 \\ &=\; -2l(I_{\beta_-} - I_{\beta_+})\frac{\rho^2}{\nu}\gamma^\varphi_L - (I_{\beta_-} - I_{\beta_+})\frac{\rho^2}{\nu}\gamma^\varphi_L.\end{aligned}$$

As a consequence of the Eq. (16a), the last term gives no contribution to T^φ_φ and we arrive at the expression

$$T^\varphi_\varphi = \frac{1}{8\pi^2}\int_0^\infty \frac{ds}{s^3}e^{-sm^2}\left\{e^{-\frac{\rho^2}{2s}}\nu^2\sum_l l\left[I_{\beta_-}(\frac{\rho^2}{2s}) - I_{\beta_+}(\frac{\rho^2}{2s})\right]\right\}^{ren}$$

Thus, using the results of the last section we obtain

$$T^t_t = T_0,\ T^\varphi_\varphi = -3T_0 + T_1.$$

Therefore one has the following structure of the energy-momentum tensor

$$\langle T^\mu_\nu\rangle^{ren} = T_0\mathrm{diag}(1,1,-3,1) + T_1\mathrm{diag}(0,0,1,0), \tag{17a}$$

where

$$T_0 \;=\; \frac{\nu}{16\pi^3}\int_0^\infty \frac{ds}{s^3}e^{-sm^2}\int_\chi \frac{e^{-iz\frac{\nu-1}{2} - \frac{\rho^2}{s}\sin^2\frac{z}{2}}}{e^{-i\nu z} - 1}dz, \tag{17b}$$

$$T_1 \;=\; -\frac{\nu m^3\cos\frac{\pi\nu}{2}}{\pi^3\rho}\int_0^\infty \frac{K_1(2m\rho\cosh\frac{y}{2})}{\cosh\frac{y}{2}}\frac{\sinh\frac{y}{2}\sinh\frac{\nu y}{2}}{\cos\nu y - \cos\pi\nu}dy. \tag{17c}$$

In the massless case, $m = 0$, we get

$$\begin{aligned} T_0 &= \frac{\nu R_1}{8\pi^3\rho^4} = -\frac{(\nu^2-1)(7\nu^2+17)}{2880\pi^2\rho^4}, \\ T_1 &= 0, \end{aligned} \tag{17d}$$

and in the massive one, we have

$$T_0 = \frac{\nu m^2 \cos\frac{\pi\nu}{2}}{2\pi^3\rho^2}\int_0^\infty \frac{K_2(2m\rho\cosh\frac{y}{2})}{\cosh^2\frac{y}{2}}\frac{\sinh\frac{y}{2}\sinh\frac{\nu y}{2}}{\cos\nu y - \cos\pi\nu}dy. \tag{17e}$$

The energy-momentum tensor obtained obeys the conservation relation:

$$\nabla_\mu T^\mu_\nu = 0.$$

Indeed, the non-trivial equation for $\nu = r$ reads as

$$\partial_r T^r_r + \frac{1}{r}(T^r_r - T^\varphi_\varphi) = 0.$$

Taking into account the manifest structure of the energy-momentum tensor given by (17), above relations takes the following form

$$\partial_r T_0 + \frac{1}{r}(4T_0 - T_1) = 0.$$

It is easy to verify that this equation is valid due to well-known relation:

$$K_2'(z) = -\frac{2}{z}K_2 - K_1.$$

The term T_1 appears due to the non-zero mass of the field. It may be represented by the following form

$$T_1 = -m^2\mathrm{Tr}[\mathcal{G}^{ren}(x;x)] = m\mathrm{Tr}[\mathcal{S}^{ren}(x;x)],$$

and gives right the trace of the energy-momentum tensor

$$\langle T^\mu_\mu\rangle^{ren} = T_1.$$

For $m\rho \gg 1$, the energy-momentum tensor is exponentially small

$$T_0 \approx \frac{\nu R_1}{8\pi^{5/2}}\frac{m^4 e^{-2m\rho}}{(m\rho)^{5/2}}.$$

Therefore the energy is localized very close to the string in a radius smaller than the Compton length of the spinor particle, $\rho < m^{-1}$. For $m\rho \ll 1$ the expression for T_k is given by (17d). The plot of the ratio $T_{0(m\neq 0)}/T_{0(m=0)}$ looks very similar to the plot in Fig.1.

V. CONCLUSION

In this paper we considered the Dirac field in the space-time of an infinitely thin and straight cosmic string. We found in manifest form the Green function and heat kernel of the squared Dirac operator. We showed that the trace of the first heat kernel coefficient is minus two times the corresponding coefficient for the scalar field in accordance with Ref. [20] and found the first three coefficients in manifest form. Thus, the energy-momentum tensor of the massive spinor field was obtained in manifest form. In the massless case the tensor is expressed in terms of the second heat kernel coefficient. The mass of the field brings an additional parameter, the Compton length of the Dirac particle. Due to this parameter the energy-momentum tensor has an Yukawa type dependence over the distance and the energy density of the vacuum polarization is concentrated close to the string, in a domain such that $\rho < m^{-1}$.

Acknowledgments

NK is grateful to Departamento de Física, Universidade Federal da Paraíba, Brazil for hospitality. This work was supported in part by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), FAPESQ-Pb/CNPq(PRONEX) and by the Russian Foundation for Basic Research Grants No. 05-02-17344, No. 05-02-39023-SFNS.

[1] Aliev A N and Gal'tsov D V 1989 *Ann. Phys.* **193** 142

[2] Aliev A N, Hortacsu M and Ozdemir N 1997 *Class. Quant. Grav.* **14** 3215

[3] Allen B and Ottewill A C 1990 *Phys. Rev.* D**42** 2669

[4] Allen B, McLaughlin J G and Ottewill A C 1992 *Phys. Rev.* D**45** 4486

[5] Allen B, Kay B S and Ottewill A C 1996 *Phys. Rev.* D**53** 6829

[6] Audretsch J, Jasper U and Skarzhinsky V D 1994 *Phys. Rev.* D**49** 6576

[7] Bordag M, Kirsten K and Dowker S 1996 *Commun. Math. Phys.* **182** 371

[8] Cheeger J 1983 *J. Diff. Geom.* **18** 575

[9] Cognola G, Kirsten K and Vanzo L 1994 *Phys. Rev.* D**49** 1029

[10] Davies P C and Sahni 1988 V *Class. Quant. Grav.* **5** 1

[11] De Lorenci V A and Moreira E S Jr 2002 *Phys. Rev.* D**65** 107503

[12] De Sousa Gerbert P and Jackiw R 1989 *Commun. Math. Phys.* **124** 229

[13] Deutsch D and Candelas P 1979 *Phys. Rev.* D**20** 3063

[14] Dowker J S 1987 *Phys. Rev.* D**36** 3095

[15] Dowker J S, Apps J S, Kirsten K and Bordag M 1996 *Class. Quantum Grav.* **13** 2911

[16] Ford L H and Vilenkin A 1981 *J. Phys.* A**14** 2353; Bezerra V B 1987 *Phys. Rev.* D**35** 2031; *id.* 1989 *J. Math. Phys.* **30** 2895

[17] Frolov V P and Serebriany E M 1987 *Phys. Rev.* D**35** 3779

[18] Frolov V P, Pinzul A and Zelnikov A I 1995 *Phys. Rev.* D**51** 2770

[19] Fursaev D V 1994 *Class. Quant. Grav.* **11** 1431

[20] Fursaev D V and Miele G 1997 *Nucl. Rev.* B**484** 697

[21] Gott J R 1985 *Astrophys. J.* **288** 422

[22] Groves P B, Anderson P R and Carlson E D 2002 *Phys.Rev.* D**66** 124017

[23] Guimarães M E X and Linet B 1994 *Commun. Math. Phys.* **165** 297

[24] Guimarães M E X 1995 *Class. Quant. Grav.* **12** 1705

[25] Helliwell T M and Konkowski D A 1986 *Phys. Rev.* D**34** 1918

[26] Harari D D and Skarzhinsky V D 1990 *Phys. Lett.* B**240** 330

[27] Iellici D 1997 *Class. Quant. Grav.* **14** 3287

[28] Iellici D and Moreira E S (Jr) 1999 *Phys. Rev.* D**60** 124015

[29] Kay B S and Studer U M 1991 *Comm. Math. Phys.* **139** 103

[30] Khusnutdinov N R 1995 *Theor. Math. Phys.* **103** 603

[31] Kibble T W B 1976 *J. Phys* A**9** 1387; Zel'dovich Ya B 1980 *Not. R. Astron. Soc.* **192** 663

[32] Kirsten K 2001 *Spectral functions in mathematics and physics* (Chapman & Hall/CRC Press)

[33] Kirsten K, Loya P and Park J 2005 *Exotic expansions and pathological properties of zeta-functions on conic manifolds*, e-Print Archive: math.sp/0511185

[34] Linet B 1986 *Phys. Rev.* D**33** 1833

[35] Linet B 1987 *Phys. Rev.* D**35** 536

[36] Linet B 1992 *Class. Quant. Grav.* **9** 2429

[37] Linet B 1995 *J. Math. Phys.* **36** 3694

[38] Linet B 1996 *Class. Quant. Grav.* **13** 97

[39] Matsas G E A 1990 *Phys. Rev.* D**42** 2927

[40] Sazhin M. *et al*, Mon. Not. R. Astron. Soc. **343**, 353 (2003)

[41] Skarzhinsky V D, Harari D D and Jasper U 1994 *Phys. Rev.* D**49** 755

[42] Smith A G 1990 *The Formation and Evolution of Cosmic Strings* ed G W Gibbons, S W Hawking and T Vachaspati (Cambridge: Cambridge University Press)

[43] Sokolov D D and Starobinsky A A 1977 *Sov. Phys. Dokl.* **22** 312

[44] Souradeep T and Sahni V 1992 *Phys. Rev.* **D46** 1616

[45] Spinelly J and Bezerra de Mello E R 2003 *Class. Quant. Grav.* **20** 873; *id.* 2004 *Int. J. Mod. Phys.* D**13** 607

[46] Vassilevich D V 2003 *Phys. Rep.* **388** 278

[47] Vilenkin A 1981 *Phys. Rev.* D**23** 852

[48] Vilenkin A and Shellard E P S 1994 *Cosmic String and other Topological Defects* (Cambridge: Cambtidge University Press)

[49] Zerbini S, Cognola G and Vanzo L 1996 *Phys. Rev.* D**54** 2669